Support Information Appendix

Hamilton's inclusive fitness maintains heritable altruism polymorphism through rb=c

Changcao Wang, and Xin Lu

Breeding social organizations across the animal kingdom: A perspective of altruism polymorphism

Reproductive altruism, with its critical influence on organisms' fitness, is the most prevalent and important form of altruistic behavior. As we have shown in the main text, the three evolutionary states of altruistic vs. non-altruistic genotypes — stable equilibrium, increasing or decreasing in frequency, which are respectively mediated by rb = c, rb > c and rb < c — all can be found from breeding social organizations across the animal kingdom. Here, based on this framework, we arrange both invertebrates and vertebrates along a continuum of social organizations in terms of altruism polymorphism. It is worth pointing out that this framework based on the rb vs. c relation is theoretical one, although in empirical studies the potential factors such as individual quality and ecological condition that affect the relation, and the potential mechanisms such as selection-mutation balance and negative frequency-dependent selection underlying the social polymorphisms, should be taken into account. We also compile information on inter-specific and intra-specific variation in breeding social organization. Furthermore, we clarify the conceptual difference between phenotype-based and genotype-based altruism polymorphism in the same population.

Our primary objective is to further illuminate why our synthesis can serve as a general guiding framework for understanding the evolution and maintenance of altruism in particular and sociality in general.

Breeding Social Organizations in the Animal Kingdom: Definition and

Distribution. Levels of sociality associated with reproduction have been measured with diverse criteria and terminology (1, 2, 3, 4). Here we define breeding social organization from a perspective of altruism polymorphism at the population scale. First, we classify breeding units within the same population of a single species as solitary and cooperative. The former refers to a unit in which one or both of the pair members are solely responsible for rearing their own offspring, and the latter refers to a unit in which individuals additional to the pair members are involved in rearing offspring of the pair. By this definition, reproductive aggregations in which there is no alloparental care (e.g. colony-nesting waterfowl on island habitats) are excluded. Then, according to differential combinations of the two types of breeding units, we can classify breeding social organizations as solitary, facultative and eusocial, and arrange

species or populations into one of the three types (Table S1, Figs. S1 and S2).

Solitary: All breeding units in a population are solitary. This type of breeding social organization is found in the majority of both invertebrate and vertebrate species. Such populations are more likely to be in monomorphic equilibrium and contain no altruistic genotype, which can be characterized by rb < c.

Facultatively cooperative (called as primitively eusocial in insects): Both solitarily and cooperatively breeding units coexist within the same population of a species. In such species, helpers (workers in insects) are usually close relatives of helped breeders. Helpers retain the potential ability to reproduce on their own, and do not differ from solitary or helped breeders in morphology, although they are often smaller and younger than the latter. Nevertheless, in a few tropical insects, for example the paper wasp *Ropalidia marginata*, cooperative groups contain helpers that stay at their natal nests and never become a queen for all of their lives (5).

This type of social organizations is exhibited by some insects and vertebrates. Cooperative groups often have higher productivity than independent breeders within the same population due to the presence of helpers. There is growing evidence that facultatively cooperative breeding systems found in insects are analogous to those in vertebrates in terms of kin-directed helping, ecological constraints as selective pressures on independent breeding, helper effect and inheritance of breeding position (6). Some facultatively cooperative breeding species with special reference to relative proportions of the two types of breeding units in the same population are listed in Table S2.

The coexistence of both types of breeding units indicates that the populations may be characterized by rb = c. Nevertheless, inconsistence between the expression of breeding social organizations and altruism polymorphism may occur on a few occasions, which will be clarified later.

Eusocial: Breeding units in a population are composed entirely of cooperative groups, known as colonies. As the extreme form of cooperative breeding, a colony often includes one or a few reproductively-active queens or kings and thousands to millions of workers, which as close relatives of the breeders have lost reproductive options for their lifetime. Therefore, a colony is usually very large compared to that of facultatively cooperative insects and vertebrates (usually less than 10 individuals), though some species such as the beetle *Austroplatypus incompertus* have colony size of 6-7 individuals. A colony also has much higher productivity than a cooperative group of facultatively cooperative breeders. Breeders and workers (called castes) in a eusocial colony are morphologically distinct, with the latter being females in some taxa or both sexes in others. Workers assist in brood care and other tasks involved in colony maintenance, such as feeding the breeders and other types of workers (7).

Eusociality is restricted to a few taxonomic groups, largely the class Insecta. The social form has also been reported in some trematode flatworms, spiders, and snapping shrimps. The naked mole rat *Heterocephalus glaber* is thought to be the only eusocial mammal.

A eusocial population is obviously altruism-monomorphic, because altruistic genotypes have gone to fixation so that they are present in all colonies in a population.

Solitarily breeding is not expected to invade the population, thereby making rb > c.

Variation in Breeding Social Organization among Taxa. While many taxonomic groups comprise only solitarily breeding species, some other groups such as ants, stingless bees, honey bees and termites include only eusocial species, because their closely related solitary taxa have gone extinct. In contrast, many lineages exhibit partial or full-range variation in the level of sociality. Those phylogenetic patterns reflect evolutionary transitions in sociality through time. For example, eusociality has 3 origins and 12 losses within the subfamily Halictinae (8). Facultatively cooperative breeding has evolved 33 times in 267 avian species (9) and has been lost 82 times in 4707 avian species (10).

Differences in social organization even occur among populations of a single species. It has been known that there is a decrease in colony size but an increase in the proportion of solitary breeders towards high latitudes or altitudes for many spider and insect species (11). In the carrion crow *Corvus corone*, all pairs in the populations of European mainland breed alone, whereas 75% of pairs in the Spanish population have helpers (12).

Potential drivers of the inter- and intra-specific variation in social organization have been attributed to environmental conditions. For short-lived insects, cold climates allow a narrow breeding window during which all individuals in a population are single-brooded, making it impossible for workers to develop (11). In facultatively cooperative breeding insects, there are at least two broods during an annual cycle, which allows the offspring from the earlier broods to act as workers helping rear the following broods. This fact may explain why eusocial lineages mostly live in the tropics. In birds, slow population turnover in warm regions is more likely to prevent some adults from finding breeding vacancies and promote them to join a pair and act as helpers; in contrast, mate shortage as the limitation on independent breeding tends to arise in cold regions where breeding turnover is fast but adult sexual ratios are male-biased (13).

Variability in social organization provides ample testing grounds for our framework in terms of altruism polymorphism based on *rb* vs. *c* relations. The evolution and maintenance of sociality can be understood via exploring internal and external factors that affect *r*, *b* and *c*. Compared to previous approaches at the between-species (review in 14) or within-colony scale (15), our framework at the within-population scale may add a new dimension to investigations of social evolution.

Conceptual Difference between Phenotype-based and Genotype-based Polymorphism in Reproductive Altruism. Obviously, solitary and eusocial taxa contain only non-altruism and altruism monomorphism, which correspond well to rb < c and rb > c, respectively. We may expect that altruism polymorphism mediated by rb = c should exist in facultatively cooperative breeding systems. Nevertheless, in some particular cases, facultatively cooperative breeding does not necessarily predict the presence of altruism polymorphism. For instance, in a few avian cooperative breeders (e.g. white-winged choughs $Corcorax\ melanorhamphus$, 16), all breeding units are cooperative but some individuals never help during their life. In contrast, it is

likely that all individuals in a population act as helpers but the population contains both solitary and cooperative nests. The inconsistency may be attributed to the difference in scales on which the two concepts rest. Altruism polymorphism by definition means that altruistic and non-altruistic genotypes are both present in a population, thereby being based on an individual scale, whereas social organization is on the breeding unit scale.

Hence, the idea of altruism polymorphism is a simple, convenient tool by which we can analyze the evolution of altruism hidden in various types of breeding social organizations.

An overview of other forms of altruism in nature: A perspective of behavioral polymorphism

In addition to reproductive altruism, a variety of altruistic behaviors have been identified in the animal kingdom. Here, we refer to altruistic behaviors as ones that reduce actors' own fitness to increase the fitness of others. Thus, we do not include reciprocal altruism in which direct benefits are exchanged between the actors and the recipients at the same time or over a period of time (17).

We provide a preliminary summary of cases of these altruistic behaviors reported in the literature (Table S3). These forms of altruism tend to occur between kin, and prove to be facultative with alternative traits coexisting in the same population. In some cases, altruism is directed towards non-kin (18, 19, 20, 21). These non-kin interactions may also be understood from the perspective of behavioral polymorphism because the terms r, b, and c in Hamilton's rule could all be positive, negative or zero (22).

- 1. Costa JT, Fitzgerald TD (1996) Developments in social terminology: semantic battles in a conceptual war. *Trends Ecol Evol* 11: 285–289.
- 2. Costa JT, Fitzgerald TD (2005) Social terminology revisited: Where are we ten years later? *Ann Zool Fenn* 42: 559–564.
- 3. Schwarz MP, Richards MH, Danforth BN (2007) Changing paradigms in insect social evolution: Insights from halictine and allodapine bees. *Annu Rev Entomol* 52: 127–150.
- 4. Dew RM, Tierney SM, Schwarz MP (2016) Social evolution and casteless societies: needs for new terminology and a new evolutionary focus. *Insec Soc* 63: 5–14.
- 5. Gadagkar R (2001) *The Social Biology of Ropalidia marginata: Toward Understanding the Evolution of Eusociality* (Cambridge Univ Press, Cambridge).
- 6. Brockmann HJ (1997) Cooperative breeding in wasps and vertebrates: the role of ecological constraints. *The Evolution of Social Behavior in Insects and Arachnids*, eds Choe JC, Crespi BJ (Cambridge Univ Press, Cambridge), pp 347–371.
- 7. Korb J, Buschmann M, Schafberg S, Liebig J, Bagnères AG (2012) Brood care and social evolution in termites. *Proc R Soc B* 279: 2662–2671.
- 8. Danforth BN (2002) Evolution of sociality in a primitively eusocial lineage of bees. *Proc Natl Acad Sci USA* 99: 286–290.
- 9. Cornwallis CK, West SA, Davis KE, Griffin AS (2010) Promiscuity and the

- evolutionary transition to complex societies. Nature 466: 969–972.
- 10. Cornwallis CK, et al. (2017) Cooperation facilitates the colonization of harsh environments. *Nat Ecol Evol* 1: 0057.
- 11. Purcell J (2011) Geographic patterns in the distribution of social systems in terrestrial arthropods. *Biol Rev* 86: 475–491.
- 12. Baglione V, et al. (2005) Does year-round territoriality rather than habitat saturation explain delayed natal dispersal and cooperative breeding in the carrion crow? *J Anim Ecol* 74: 842–851.
- 13. Zhang GY, Zhao QT, Møller AP, Komdeur J, Lu X (2017) Climate predicts which sex acts as helpers among cooperatively breeding bird species. *Biol Lett* 13: 20160863.
- 14. Bourke AF (2014) Hamilton's rule and the causes of social evolution. *Phil Trans R Soc B* 369: 20130362.
- 15. Rehan SM, Toth AL (2015) Climbing the social ladder: The molecular evolution of sociality. *Trends Ecol Evol* 30: 426–433.
- 16. Rowley I (1978) Communal activities among white-winged choughs *Corcorax melanorhamphus*. *Ibis* 120: 178–197.
- 17. West SA, Griffin AS, Gardner A (2007) Social semantics: Altruism, cooperation, mutualism, strong reciprocity and group selection. *J Evol Biol* 20: 415–432.
- 18. McDonald DB, Potts WK (1994) Cooperative display and relatedness among males in a lek-mating bird. *Science* 266: 1030–1032.
- 19. Loiselle BA, et al. (2006) Kin selection does not explain male aggregation at leks of 4 manakin species. *Behav Ecol* 18: 287–291.
- 20. DuVal EH (2007) Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. *Am Nat* 169: 423–432.
- 21. Riehl C (2013) Evolutionary routes to non-kin cooperative breeding in birds. *Proc R Soc Lond B* 280: 20132245.
- 22. Gardner A, West SA, Wild G (2011) The genetical theory of kin selection. *J Evol Biol* 24: 1020–1043.




Fig. S1. A sketch showing different types of breeding social organization. a. Solitary: a population consisting completely of solitarily breeding units; b. Facultatively cooperative: a population consisting of both solitarily and cooperatively breeding units; c. Eusocial: a population consisting completely of cooperatively breeding units.



Fig. S2. Making sense of altruism coexisting with non-altruism in nature and humans. Helping to rear offspring of others in facultatively cooperative breeders: a. paper wasps *Polistes dominulus*; b. Tibetan ground tits; c. meerkats *Suricata suricatta*. Several other forms of altruism: d. Adaptive suicide, to reduce the chance of kin being parasitized in pea aphids *Acyrthosiphon pisum*; e. Cooperative courtship, to facilitate mating success of other males in wild turkeys *Meleagris galloparvo*; f. Compassion for others, to give alms to beggars in humans. Photos a-e by Eugene Zelenko, Danhua Ke, Charles Sharp, Shipher Wu, and Kelly Burgess, respectively, and the painting f by Jacques-Louis David.

Table S1. The distribution of breeding social organizations across the animal kingdom

| Class | Order | | No. of | | | e Eusocial |
|----------------------------------|----------------------|------------------------------|------------|----------|-------------|------------|
| | | describe Family or subfamily | described | | Facultative | |
| | | ranning or subtaining | species in | Solitary | | Eusociai |
| | | | the world | | | |
| Invertebrates | | | | | | |
| Digenea (trematode) ¹ | | | 6000 | most | most | a few |
| Malacostraca ² | Decapoda | Alpheidae (shrimps) | 1120 | most | a few | a few |
| Arachnida ³ | Araneae (spiders) | | 40700 | most | a few | a few |
| Insecta ^{4, 5, 6, 7, 8} | Hymenoptera | Formicidae (ants) | 16000 | 0 | 0 | all |
| | | Apidae (bees) | 5700 | many | many | many |
| | | Vespidae (wasps) | 5000 | many | many | many |
| | | Crabronidae (wasps) | 9000 | many | a few | 0 |
| | | Halictinae (bees) | 2500 | many | 830 | 0 |
| | | Apidae (bees) | 5700 | many | a few | a few |
| | Isoptera (termites) | | 2600 | a few | a few | most |
| | Hemiptera | Aphididae (aphids) | 2280 | most | a few | a few |
| | Thysanoptera | Thripidae (thrips) | 2000 | most | a few | a few |
| | Coleoptera (beetles) | | 400000 | most | 1 | 1 |
| Vertebrates | | | | | | |
| Osteichthyes ⁹ | Perciformes | Cichlaidae (cichlids) | 2000 | most | a few | 0 |
| Aves (birds) 10 | | | 10000 | most | 900 | 0 |
| Mammalia (mammals) 11 | | | 5420 | most | 100 | 1 |

1. Hechinger *et al.* 2011; 2. Duffy & Thie 2007; 3. Avil & 1997; 4. Wilson 1971; 5. Kent & Simpson 1992; 6. Schwarz *et al.* 2007; 7. Korb *et al.* 2012; 8. Rehan & Toth 2015; 9. Wong & Balshine 2011; 10. Cockburn 2006; 11. Burda *et al.* 2000.

Avil & L (1997) Causes and consequences of cooperation and permanent-sociality in spiders. *The Evolution of Social Behavior in Insects and Arachnids*, eds Choe JC, Crespi BJ (Cambridge Univ Press, Cambridge), pp 476–498.

Burda H, Honeycutt RL, Begall S, Locker-Grütjen O, Scharff A (2000) Are naked and common mole-rats eusocial and if so, why? *Behav Ecol Sociobiol* 47: 293–303.

Cockburn A (2006) Prevalence of different modes of parental care in birds. *Proc R Soc B* 273: 1375–1383.

Duffy JE, Thie M (2007) Sexual and social behavior of crustacea. *Evolutionary Ecology of Social and Sexual Systems: Crustaceans as Model Organisms*, eds Duffy JE, Thiel M (Oxford Univ Press, Oxford), pp 387–409.

Hechinger RF, Wood AC, Kuris AM (2011) Social organization in a flatworm: trematode parasites form soldier and reproductive castes. *Proc R Soc Lond B* 278: 656–665.

Kent DS, Simpson JA (1992) Eusociality in the beetle Austroplatypus incompertus

- (Coleoptera: Curculionidae). Naturwissenschaften 79: 86-87.
- Korb J, Buschmann M, Schafberg S, Liebig J, Bagnères AG (2012) Brood care and social evolution in termites. *Proc R Soc B* 279: 2662–2671.
- Rehan SM, Toth AL (2015) Climbing the social ladder: The molecular evolution of sociality. *Trends Ecol Evol* 30: 426–433.
- Schwarz MP, Richards MH, Danforth BN (2007) Changing paradigms in insect social evolution: Insights from halictine and allodapine bees. *Annu Rev Entomol* 52: 127–150.
- Wilson EO (1971) The Insect Societies (Harvard Univ Press, Cambridge).
- Wong M, Balshine S (2011) The evolution of cooperative breeding in the African cichlid fish, *Neolamprologus pulcher*. *Biol Rev* 86: 511–530.

Table S2. A summary showing the relative proportion of solitarily and cooperatively breeding units in a population of facultatively cooperative breeding insects and vertebrates

| Scientific name | No. of | % cooperatively | Courage | |
|-------------------------|----------------|-----------------|-------------------------------------|--|
| Scientific name | breeding units | breeding unit | Sources | |
| Insects | | | | |
| Myrmecocystus mimicus | 96 | 53.1 | Bartz & Holldobler 1982 | |
| Polistes fuscatus | 38 | 2.6 | Eberhard 1969 | |
| Polistes canadensis | 20 | 25.0 | Eberhard 1969 | |
| Xylocopa virginica | 23 | 65.2 | Prager 2014 | |
| Ropalidia marginata | 48 | 77.1 | Premnath et al. 1996 | |
| Ceratina nigrolateralis | 32 | 6.7 | Rehan et al. 2009 | |
| Neoceratina dentipes | 20 | 5.0 | Rehan et al. 2009 | |
| Pithitis smaragdula | 5 | 20.0 | Rehan et al. 2009 | |
| Ceratina australensis | 378 | 12.4 | Rehan et al. 2011 | |
| Lasioglossum malachurum | 52 | 21.2 | Richards et al.2005 | |
| Ceratina japonica | | 20.0 | Sakagami & Maeta 1984 | |
| Megalopta genalis | 113 | 65.5 | Kapheim et al. 2013 | |
| Polistes metricus | 64 | 21.9 | Metcalf & Whitt 1977 | |
| Lasius niger | | 18.0 | Sommer & Holldobler 1995 | |
| Xylocopa sulcatipes | 124 | 43.5 | Stark 1992 | |
| Exoneura nigrescens | 46 | 19.6 | Zammit et al. 2008 | |
| Fish | | | | |
| Neolamprologus pulcher | 60 | 95.0 | Taborsky & Limberger 1981, Taborsky | |
| Birds | | | | |
| Rhea americana | 35 | 23.0 | Codenotti & Alvarez 1997 | |
| Tetraophasis szechenyii | 68 | 64.7 | Xu et al. 2011 | |
| Opisthocomus hoazin | 364 | 59.0 | Strahl 1988 | |
| Gypaetus barbatus | 92 | 15.2 | Bertran & Margalida 2002 | |
| Haliaeetus vociferoides | 95 | 42.0 | Tingay et al. 2002 | |
| Neophron percnopterus | 37 | 5.4 | Tella 1993 | |
| Parabuteo unicinctus | 64 | 84.4 | James & Mannan 1991 | |
| Buteo galapagoensis | 32 | 68.8 | Faaborg et al. 1980 | |
| Melierax canorus | 117 | 13.7 | Malan 2004 | |
| Monias benschi | 108 | 56.5 | Seddon et al. 2003 | |
| Porphyrio porphyrio | 93 | 92.0 | Craig & Jamieson 1990 | |
| Psophia leucoptera | 88 | 100.0 | Sherman 1995 | |
| Rhynochetos jubatus | 38 | 89.5 | Theuerkauf et al. 2009 | |

| Gallinula chloropus | 97 | 12.4 | Gibbons 1986 |
|---------------------------|-----|-------|---------------------------------|
| Gallinula mortierii | 106 | 54.7 | Goldizen et al. 1998 |
| Vanellus chilensis | 23 | 47.8 | Saracura et al. 2008 |
| Eclectus roratus | 34 | 58.8 | Heinsohn & Legge 2003 |
| Myiopsitta monachus | 15 | 20.0 | Eberhard 1998 |
| Pyrrhura orcesi | 35 | 74.0 | Ridgely & Robbins 1988 |
| Coracopsis vasa | 15 | 100.0 | Ekstrom et al. 2007 |
| Crotophaga sulcirostris | 171 | 76.0 | Koford et al. 1990 |
| Guira guira | 39 | 71.8 | Lima et al. 2011 |
| Crotophaga ani | 71 | 100.0 | Grieves et al. 2014 |
| Crotophaga major | 87 | 8.0 | Riehl & Jara 2009 |
| Ceryle rudis | 59 | 45.8 | Reyer 1990 |
| Dacelo novaeguineae | 131 | 66.7 | Legge & Cockburn 2000 |
| Merops apiaster | 239 | 19.0 | Lessells 1990 |
| Merops bullockoides | 187 | 49.7 | Emlen & Wrege 1988 |
| Merops bulocki | 96 | 28.1 | Fry 1972 |
| Merops leschenaulti | 3 | 33.3 | Burt 2002 |
| Merops orientalis | 24 | 37.5 | Sridhar & Karanth 1993 |
| Merops ornatus | 180 | 45.0 | Boland 2004 |
| Merops philippinus | 19 | 42.0 | Burt 2002 |
| Penelopides exarhatus | 2 | 100.0 | O'brien 1997 |
| Phoeniculus damarensis | 22 | 100.0 | Plessis et al. 2007 |
| Phoeniculus purpureus | 189 | 90.5 | Ligon 1990 |
| Todiramphus cinnamominus | 57 | 54.5 | Kesler & Haig 2007 |
| Tanysiptera sylvia | 71 | 7.0 | Legge & Heinsohn 2001 |
| Colaptes campestris | 26 | 42.0 | Dias et al. 2013 |
| Semnornis ramphastinus | 28 | 62.0 | Restrepo & Mondrag ón 1998 |
| Picoides borealis | 93 | 46.0 | Lennartz et al. 1987 |
| Acanthisitta chloris | 105 | 10.5 | Sherley 1990 |
| Acrocephalus sechellensis | 657 | 29.0 | Richardson et al. 2002 |
| Acrocephalus vaughani | 22 | 36.0 | Brooke & Hartley 1995 |
| Alophoixus pallidus | 183 | 60.1 | Sankamethawee et al. 2010, 2011 |
| Aphelocoma coerulescens | 545 | 55.0 | Woolfenden & Fitzpatrick 1990 |
| Babax waddelli | 10 | 40.0 | Lu 2004 |
| Bubalornis niger | 137 | 100.0 | Winterbottom et al. 2001 |
| Calocitta formosa | 56 | 52.0 | Innes & Johnston 1996 |
| Campylorhynchus griseus | 90 | 30.0 | Austad & Rabenold 1986 |
| Campylorhynchus nuchalis | 230 | 67.4 | Rabenold 1990 |
| Climacteris affinis | 26 | 34.6 | Radford 2004 |
| Climacteris erythrops | 20 | 65.0 | Noske 1991 |
| | | | |

| Climacteris picumnus | 102 | 38.2 | Doerr & Doerr 2006 |
|---------------------------|-----|-------|--------------------------|
| Climacteris rufus | 90 | 59.0 | Luck 2001 |
| Corcorax melanorhamphos | 186 | 98.9 | Rowley 1978 |
| Corvus brachyrhynchos | 115 | 30.0 | Caffrey 1992 |
| Corvus corone | 236 | 75.0 | Baglione et al. 2002 |
| Cyanocorax beecheii | 36 | 81.0 | Raitt et al. 1984 |
| Cyanocorax chrysops | 6 | 100.0 | Uejima et al. 2012 |
| Cyanocorax morio | 46 | 100.0 | Williams & Rabenold 2005 |
| Cyanopica cyanus | 110 | 49.1 | Valencia et al. 2003 |
| Eopsaltria georgiana | 258 | 70.0 | Russell et al. 2004 |
| Erythropygia coryphaeus | 427 | 15.0 | Lloyd et al. 2009 |
| Garrulax perspicillatus | 19 | 100.0 | Wang 2011 |
| Gymnorhina tibicen | 12 | 100.0 | Finn & Hughes 2001 |
| Gymnorhinus | 341 | 8.0 | Marzluff & Balda 1990 |
| Lamprotornis pulcher | 36 | 97.2 | Wilkinson 1982 |
| Lamprotornis superbus | 352 | 91.0 | Rubenstein 2007 |
| Lanius excubitoroides | 82 | 81.7 | Zack 1986 |
| Malurus coronatus | 73 | 46.6 | Kingma et al. 2010 |
| Malurus cyaneus | 63 | 36.5 | Nias & Ford 1992 |
| Malurus elegans | 398 | 83.0 | Russell & Rowley 2000 |
| Malurus leucopterus | 37 | 48.6 | Rowley & Russell 1995 |
| Malurus melanocephalus | 604 | 20.0 | Varian-Ramos et al. 2010 |
| Malurus splendens | 226 | 65.0 | Russell & Rowley 1988 |
| Manorina melanophrys | 31 | 100.0 | Clarke et al. 2002 |
| Melanerpes formicivorus | 354 | 76.5 | Koenig & Stacey 1990 |
| Melanerpes formicivorus | 164 | 40.2 | Koenig & Stacey 1990 |
| Melanerpes formicivorus | 32 | 14.6 | Koenig & Stacey 1990 |
| Mimus parvulus | 450 | 34.0 | Curry & Grant 1989 |
| Myrmecocichla formicivora | 87 | 74.7 | Barnaby 2012 |
| Neothraupis fasciata | 71 | 32.4 | Manica & Marini 2012 |
| Philetairus socius | 94 | 58.5 | Covas et al. 2006 |
| Phyllastrephus cabanisi | 103 | 37.9 | Callens 2012 |
| Pomatostomus ruficeps | 90 | 94.0 | Browning et al. 2012 |
| Pomatostomus temporalis | 75 | 86.7 | Eguchi et al. 2007 |
| Prunella collaris | 96 | 44.8 | Nakamura 1998 |
| Prunella modularis | 208 | 70.2 | Davies 1986 |
| Psaltriparus minimus | 97 | 35.0 | Sloane 1996 |
| Pseudonigrita arnaudi | 68 | 23.9 | Bennun 1994 |
| Pseudopodoces humilis | 159 | 28.0 | Lu et al. 2011 |
| Schetba rufa | 220 | 37.3 | Eguchi et al. 2002 |

| Sericornis frontalis | 169 | 54.0 | Magrath & Whittingham 1997 |
|--|--|---|---|
| Sialia mexicana | 741 | 7.4 | Dickinson et al. 1996 |
| Struthidea cinerea | 61 | 100.0 | Woxvold et al. 2006 |
| Stipiturus malachurus | 48 | 8.0 | Maguire & Mulder 2004 |
| Lichenostomus melanops | 7 | 28.6 | Franklin et al. 1995 |
| Manorina melanocephala | 35 | 100.0 | Poldmaa et al. 1995 |
| Notiomystis cincta | 30 | 26.7 | Castro et al. 1996 |
| Daphoenositta chrysoptera | 28 | 82.1 | Noske 1998 |
| Chaetops frenatus | 35 | 34.3 | Holmes et al. 2002 |
| Plocepasser mahali | 132 | 66.7 | Harrison et al. 2013 |
| Ramphocinclus brachyurus | 112 | 34.8 | Temple et al. 2009 |
| Sitta pusilla | 347 | 19.7 | Cox & Slater 2007 |
| Sitta pygmaea | 141 | 39.7 | Sydeman et al.1988 |
| Turdoides bicolor | 181 | 94.5 | Ridley & Raihani 2008 |
| Turdoides caudata | 20 | 50.0 | Gaston 1978a |
| Turdoides squamiceps | 19 | 89.5 | Zahavi 1990 |
| Turdoides striata | 30 | 100.0 | Gaston 1978b |
| Yuhina brunneiceps | 69 | 97.1 | Yuan et al. 2004 |
| Malurus pulcherrimus | 121 | 27.0 | Rowley & Russell 2002 |
| Corvinella corvina | 18 | 100.0 | Grimes 1980 |
| Eopsaltria australis | 29 | 27.6 | Debus 2006 |
| Cisticola chiniana | 17 | 88.2 | Carlson 1986 |
| Acrocephalus melanopogon | 25 | 68.0 | Blomqvist et al. 2005 |
| Acrocephalus baeticatus | 65 | 12.0 | Eising et al. 2001 |
| Mimus macdonaldi | 53 | 67.9 | Lippke 2008 |
| Catharus bicknelli | 18 | 78.0 | Goetz et al. 2003 |
| Loxioides bailleui | 25 | 24.0 | Patch-Highfill 2008 |
| Calcarius pictus | 23 | 82.6 | Briskie et al. 1998 |
| Hypopyrrhus | 27 | 100.0 | Ocampo et al. 2012 |
| Agelaioides badius | 20 | 18.0 | Fraga 1991 |
| Parus fasciiventer | 49 | 61.0 | Shaw et al. 2015 |
| Aegithalos glaucogularis | 21 | 42.9 | Li et al. 2014 |
| Aegithalos concinnus | 50 | 20.0 | Li et al. 2012 |
| Garrulax perspicillatus | 24 | 79.2 | Wang 2011 |
| Mammals | | | |
| | 211 | 84.4 | Dietz & Baker1993 |
| - | | | |
| _ | | | • |
| | | | |
| Acrocephalus melanopogon Acrocephalus baeticatus Mimus macdonaldi Catharus bicknelli Loxioides bailleui Calcarius pictus Hypopyrrhus Agelaioides badius Parus fasciiventer Aegithalos glaucogularis Aegithalos concinnus | 25 65 53 18 25 23 27 20 49 21 | 68.0 12.0 67.9 78.0 24.0 82.6 100.0 18.0 61.0 42.9 20.0 | Blomqvist et al. 2005 Eising et al. 2001 Lippke 2008 Goetz et al. 2003 Patch-Highfill 2008 Briskie et al. 1998 Ocampo et al. 2012 Fraga 1991 Shaw et al. 2015 Li et al. 2014 Li et al. 2012 |

| 7 | 57.1 | Spinks et al. 2000 |
|-----|---|--|
| 6 | 100.0 | Scharff et al. 2000 |
| 28 | 53.6 | Yeboah & Dakwa 2002 |
| 1 | 100.0 | Jarvis 1981 |
| 14 | 85.7 | Schradin & Pillay 2004 |
| 9 | 44.4 | Agren et al. 1989 |
| 243 | 15.6 | Getz & Hofmann 1986 |
| 20 | 65.0 | FitzGerald & Madison 1983 |
| 27 | 29.6 | Wolff 1994 |
| 848 | 7.1 | Smith 1966 |
| 13 | 46.2 | Strand et al. 2000 |
| 19 | 57.9 | Moehlman 1987 |
| 8 | 62.5 | Bekoff & Wells 1986 |
| 24 | 62.5 | Harrington et al. 1983 |
| 15 | 73.3 | Moehlman 1979 |
| 61 | 90.2 | Zubiri & Gottelli 1995 |
| 170 | 99.0 | Rood 1990 |
| 26 | 100.0 | Fuller et al. 1992 |
| 9 | 100.0 | Doolan & Mcdonald 1997 |
| 14 | 100.0 | Cant 2000 |
| | 6 28 1 14 9 243 20 27 848 13 19 8 24 15 61 170 26 | 6 100.0 28 53.6 1 100.0 14 85.7 9 44.4 243 15.6 20 65.0 27 29.6 848 7.1 13 46.2 19 57.9 8 62.5 24 62.5 15 73.3 61 90.2 170 99.0 26 100.0 |

Insects

- Bartz SH, Hölldobler B (1982) Colony founding in *Myrmecocystus mimicus* Wheeler (Hymenoptera: Formicidae) and the evolution of foundress associations. *Behav Ecol Sociobiol* 10: 137–147.
- Eberhard MJW (1969)The social biology of polistine wasps. *Misc Publ Mus Zool Univ Mich* 140: 1–101.
- Kapheim KM, Smith AR, Nonacs P, Wcislo WT, Wayne RK (2013) Foundress polyphenism and the origins of eusociality in a facultatively eusocial sweat bee, *Megalopta genalis* (Halictidae). *Behav Ecol Sociobiol* 67: 331–340.
- Metcalf RA, Whitt GS (1977) Relative inclusive fitness in the social wasp *Polistes metricus*. *Behav Ecol Sociobiol* 2: 353–360.
- Prager SM (2014) Comparison of social and solitary nesting carpenter bees in sympatry reveals no advantage to social nesting. *Biol J Linnean Soc* 113: 998–1010.
- Premnath S, Sinha A, Gadagkar R (1996) Dominance relationship in the establishment of reproductive division of labour in a primitively eusocial wasp (*Ropalidia marginata*). *Behav Ecol Sociobiol* 39: 125–132.
- Rehan SM, Richards MH, Schwarz MP (2009) Evidence of social nesting in the *Ceratina* of Borneo (Hymenoptera: Apidae). *J Kans Entomol Soc* 82: 194–209.
- Rehan SM, Richards MH, Schwarz MP (2010) Social polymorphism in the Australian small carpenter bee, *Ceratina* (Neoceratina) *australensis*. *Insect Soc* 57:

- 403-412.
- Richards MH, French D, Paxton RJ (2005) It's good to be queen: classically eusocial colony structure and low worker fitness in an obligately social sweat bee. *Mol Ecol* 14: 4123–4133.
- Sakagami SF, Maeta Y (1984) Multifemale nests and rudimentary castes in the normally solitary bee *Ceratina japonica* (Hymenoptera: Xylocopinae). *J Kans Entomol Soc* 57: 639–656.
- Smith AR, Wcislo WT, O'Donnell S (2007) Survival and productivity benefits to social nesting in the sweat bee *Megalopta genalis* (Hymenoptera: Halictidae). *Behav Ecol Sociobiol* 61: 1111–1120.
- Sommer K, Hoelldobler B (1995) Colony founding by queen association and determinants of reduction in queen number in the ant *Lasius niger*. *Anim Behav* 50: 287–294.
- Stark RE (1992) Cooperative nesting in the multivoltine large carpenter bee *Xylocopa sulcatipes* Maa (Apoidea: Anthophoridae): do helpers gain or lose to solitary females? *Ethology* 91: 301–310.
- Zammit J, Hogendoorn K, Schwarz MP (2008) Strong constraints to independent nesting in a facultatively social bee: quantifying the effects of enemies-at-the-nest. *Insect Soc* 55: 74–78.

Fish

Taborsky M, Limberger D (1981) Helpers in fish. *Behav Ecol Sociobiol* 8: 143–145. Taborsky M (1984) Brood care helpers in the cichlid fish *Lamp rologus brichardi*: their costs and benefits. *Anim Behav* 32: 1236–1252.

Birds

- Austad S, Rabenold K (1986) Demography and the evolution of cooperative breeding in the bicolored wren, *Campylorhynchus griseus*. *Behaviour* 97: 308–324.
- Baglione V, Marcos J, Canestrari D (2002) Cooperatively breeding groups of the carrion crow *Corvus corone corone* in northern Spain. *Auk* 119: 790–799.
- Barnaby J (2012) Cooperative breeding in the southern anteater-chat: sexual disparity survival and dispersal. PhD thesis (Uppsala Univ, Uppsala, Sweden).
- Bennun L (1994) The contribution of helpers to feeding nestlings in grey-capped social weavers *Pseudonigrita arnaudi*. *Anim Behav* 47: 1047–1056.
- Bertran J, Margalida A (2002) A Social organization of a trio of bearded vulture (*Gypaetus barbatus*): social and parental roles. *J Raptor Res* 36: 66–70.
- Blomqvist D, Fessl B, Hoi H, Kleindorfer S (2005) High frequency of extra-pair fertilisations in the moustached warbler, a songbird with a variable breeding system. *Behaviour* 142: 1133–1148.
- Boland CRJ (2004) Breeding biology of rainbow bee-eaters (*Merops ornatus*): a migratory, colonial, cooperative bird. *Auk*121: 811–823.
- Briskie JV, Montgomerie R, P õldmaa T, Boag PT (1998) Paternity and parental care in the polygynandrous Smith's longspur. *Behav Ecol Sociobiol* 43: 181–190.
- Brooke M, Hartley IR (1995) Nesting Henderson reed-warblers (Acrocephalus

- *vaughani taiti*) studied by DNA fingerprinting: unrelated coalitions in a stable habitat? *Auk* 112: 77–86.
- Browning LE, et al. (2012) Carer provisioning rules in an obligate cooperative breeder: prey type, size and delivery rate. *Behav Ecol Sociobiol* 66: 1639–1649.
- Burt DB, Peterson AT (1993) Biology of cooperative-breeding scrub jays (*Aphelocoma coerulescens*) of Oaxaca, Mexico. *Auk* 110: 207–214.
- Burt DB (2002) Social and breeding biology of bee-eaters in Thailand. *Wilson Bull* 114: 275–279.
- Caffrey C (1999) Feeding rates and individual contributions to feeding at nests in cooperatively breeding western American crows. *Auk* 116: 836–841.
- Callens T (2012) Genetic and demographic signatures of population fragmentation in a cooperatively breeding bird from south-east Kenya. PhD thesis (Ghent Univ, Ghent, Belgium).
- Carlson A (1986) Group territoriality in the rattling cisticola, *Cisticola chiniana*. *Oikos* 47: 181–189.
- Castro I, Minot EO, Fordham RA, Birkhead TR (1996) Polygynandry, face-to-face copulation and sperm competition in the Hihi *Notiomystis cincta* (Aves: Meliphagidae). *Ibis* 138: 765–771.
- Clarke MF, et al. (2002) Male-biased sex ratios in broods of the cooperatively breeding bell miner *Manorina melanophrys*. *J Avian Biol* 33: 71–76.
- Codenotti TL, Alvarez F (1997) Cooperative breeding between males in the greater rhea *Rhea americana*. *Ibis* 139: 568–571.
- Covas R (2002) Life-history evolution and cooperative breeding in the sociable weaver. PhD thesis (Univ of Cape Town, Cape Town, South Africa).
- Cox JA, Slater GL (2007) Cooperative breeding in the brown-headed nuthatch. *Wilson J Ornithol* 119: 1–8.
- Craig JL, Jamieson IG (1990) Pukeko: Different approaches and some different answers. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 385–412.
- Curry RL, Grant PR (1989) Demography of the cooperatively breeding Galapagos mockingbird, *Nesomimus parvulus*, in a climatically variable environment. *J Anim Ecol* 58: 441–463.
- Davies NB, Houston AI (1986) Reproductive success of dunnocks, *Prunella modularis*, in a variable mating system. II. Conflicts-of-interest among breeding adults. *J Anim Ecol* 55: 139–154.
- Debus SJS (2006) Breeding and population parameters of robins in a woodland remnant in northern New South Wales Australia. *Emu* 106: 147–156.
- Dias RI, Macedo RH, Goedert D, Webster MS (2013) Cooperative breeding in the Campo flicker I: breeding ecology and social behavior. *Condor* 115: 847–854.
- Dickinson JL, Koenig WD, Pitelka FA (1996) Fitness consequences of helping behavior in the western bluebird. *Behav Ecol* 7: 168–177.
- Doerr ED, Doerr VAJ (2006) Comparative demography of treecreepers: evaluating hypotheses for the evolution and maintenance of cooperative breeding. *Anim*

- Behav 72: 147-159.
- du Plessis MA, Simmons RE, Radford AN (2007) Behavioural ecology of the Namibian Violet Woodhoopoe *Phoeniculus damarensis*. *Ostrich* 78: 1–5.
- Eberhard JR (1998) Breeding biology of the monk parakeet. *Wilson Bull* 110: 463–473.
- Eguchi K, Yamagishi S, Asai S, Nagata H, Hino T (2002) Helping does not enhance reproductive success of cooperatively breeding rufous vanga in Madagascar. *J Anim Ecol* 71: 123–130.
- Eguchi K, et al. (2007) Social structure and helping behaviour of the grey-crowned babbler *Pomatostomus temporalis*. *J Ornithol* 148: S203–S210.
- Eising CM, Komdeur J, Buys J, Reemer M, Richardson DS (2001) Islands in a desert: breeding ecology of the African reed warbler *Acrocephalus baeticatus* in Namibia. *Ibis* 143: 482–493.
- Ekstrom JMM, Burke T, Randrianaina L, Birkhead TR (2007) Unusual sex roles in a highly promiscuous parrot: the Greater Vasa parrot *Caracopsis vasa*. *Ibis* 149: 313–320.
- Emlen ST, Wrege PH (1988) The role of kinship in helping decisions among white-fronted bee-eaters. *Behav Ecol Sociobiol* 23: 305–315.
- Faaborg J, Vries TD, Patterson CB, Griffin CR (1980) Preliminary observations on the occurrence and evolution of polyandry in the Galapagos hawk (*Buteo galapagoensis*). *Auk* 97: 581–590.
- Finn PG, Hughes JM (2001) Helping behaviour in Australian magpies, *Gymnorhina tibicen*. *Emu* 101: 57–63.
- Fraga RM (1991) The social system of a communal breeder, the bay-winged cowbird *Molothrus badius*. *Ethology* 89: 195–210.
- Franklin DC, Smales IJ, Miller MA, Menkhorst PW (1995) The reproductive biology of the helmeted honeyeater, *Lichenostomus melanops cassidix*. *Wildl Res* 22: 173–191.
- Fry CH (1972) The social organization of bee-eaters (*Meropidae*) and cooperative breeding in hot-climate birds. *Ibis* 114: 1–14.
- Gaston AJ (1978a) Ecology of the common babbler *Turdozdes caudatus*. *Ibis* 120: 415–432.
- Gaston AJ (1978b) Demography of the jungle babbler *Turdoides striatus*. *J Anim Ecol* 47: 845–879.
- Gibbons DW (1986) Brood Parasitism and Cooperative Nesting in the moorhen, *Gallinula chloropus. Behav Ecol Sociobiol* 19: 221–232.
- Goetz JE, McFarland KP, Rimmer CC (2003) Multiple paternity and multiple male feeders in Bicknell's thrush (*Catharus bicknelli*). *Auk* 120: 1044–1053.
- Goldizen AW, Putland DA, Goldizen AR (1998) Variable mating patterns in Tasmanian native hens (*Gallinula mortierii*): correlates of reproductive success. *J Anim Ecol* 67: 307–317.
- Grieves LA, Logue DM, Quinn JS (2014) Joint-nesting smooth-billed anis, *Crotophaga ani*, use a functionally referential alarm call system. *Anim Behav* 89: 215–221.

- Grimes LG (1980) Observations of group behaviour and breeding biology of the yellow-billed shrike *Corvinella corvina*. *Ibis* 122: 166–192.
- Harrison XA, York JE, Cram DL, Hares MC, Young AJ (2013) Complete reproductive skew within white-browed sparrow weaver groups despite outbreeding opportunities for subordinates of both sexes. *Behav Ecol Sociobiol* 67: 1915–1929.
- Heinsohn R, Legge S (2003) Breeding biology of the reverse-dichromatic, cooperative parrot *Eclectus roratus*. *J Zool* 259: 197–208.
- Holmes RT, Frauenknecht BD, du Plessis MA (2002) Breeding system of the cape rockjumper, a South African fynbos endemic. *Condor* 104: 188–192.
- Innes KE, Johnston RE (1996) Cooperative breeding in the white-throated magpie-jay: how do auxiliaries influence nesting success? *Anim Behav* 51: 519–533.
- James WD, Mannan RW (1991) Dominance hierarchies and helper contributions in Harris' hawks. *Auk* 108: 649–660.
- Kesler DC, Haig SM (2007) Territoriality, prospecting, and dispersal in cooperatively breeding Micronesian kingfishers (*Todiramphus cinnamominus reichenbachii*). *Auk* 124: 381–395.
- Kingma SA, Hall ML, Arriero E, Peters A (2010) Multiple benefits of cooperative breeding in purple-crowned fairy-wrens: a consequence of fidelity? *J Anim Ecol* 79: 757–768.
- Koenig WD, Stacey PB (1990) Acorn woodpeckers: group-living and food storage under contrasting ecological conditions. *Cooperative Breeding in Birds:*Long-term Studies of Ecology and Behavior, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 413–454.
- Koford RR, Bowen BS, Vehrencamp SL (1990) Groove-billed anis: joint-nesting in a tropical cuckoo. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 333–355.
- Legge S, Cockburn A (2000) Social and mating system of cooperatively breeding laughing kookaburras (*Dacelo novaeguineae*). *Behav Ecol Sociobiol* 47: 220–229.
- Legge S, Heinsohn R (2001) Kingfishers in paradise: the breeding biology of *Tanysiptera sylvia* at the Iron Range National Park, Cape York. *Aust J Zool* 49: 85–98.
- Lennartz M, Hooper R, Harlow R (1987) Sociality and cooperative breeding of red-cockaded woodpeckers, *Picoides borealis*. *Behav Ecol Sociobiol* 20: 77–88.
- Lessells CM (1990) Helping at the nest in European bee-eaters: who helps and why? *Population Biology of Passerine Birds: An Integrated Approach*, eds Blondel J, Gosler A, Lebreton JD, MacCleery R (Springer-Verlag, Berlin), pp 357–368.
- Li J, Lv L, Wang Y, Xi B, Zhang Z (2012) Breeding biology of two sympatric *Aegithalos* tits with helpers at the nest. *J Ornithol* 153: 273–283.
- Li SH, Brown JL (2000) High frequency of extrapair fertilization in a plural breeding bird the Mexican jay, revealed by DNA microsatellites. *Anim Behav* 60: 867–877.

- Ligon JD, Ligon SH (1990) Green woodhoopoes: life history traits and sociality. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 31–66.
- Lima MR, Macedo RH, Muniz L, Pacheco A, Graves JA (2011) Group composition, mating system, and relatedness in the communally breeding guira cuckoo (*Guira guira*) in central Brazil. *Auk* 128: 475–486.
- Lippke V, Ilonka S (2008) Ecology and evolution of cooperation in the Española mockingbird, *Nesomimus macdonaldi*, Gal ápagos. PhD thesis (Univ of California Los Angeles, Los Angeles, CA).
- Lloyd P, Taylor WA, du Plessis MA, Martin TE (2009) Females increase reproductive investment in response to helper-mediated improvements in allo-feeding, nest survival, nestling provisioning and post-fledging survival in the Karoo scrub-robin *Cercotrichas coryphaeus*. *J Avian Biol* 40: 400–411.
- Lu X (2004) Conservation status and reproductive ecology of giant babax *Babax* waddelli (Aves, Timaliinae) endemic to the Tibet plateau. *Oryx* 38: 418–425.
- Lu X, Huo R, Li Y, Liao W, Wang C (2011) Breeding ecology of ground tits in northeastern Tibetan plateau, with special reference to cooperative breeding system. *Curr Zool* 57: 751–757.
- Luck GW (2001) The demography and cooperative breeding behaviour of the rufous treecreeper *Climacteris rufa*. *Aust J Zool* 49: 515–537.
- Magrath RD, Whittingham LA (1997) Subordinate males are more likely to help if unrelated to the breeding female in cooperatively breeding white-browed scrubwrens. *Behav Ecol Sociobiol* 41: 185–192.
- Maguire GS, Mulder AM (2004) Breeding biology and demography of the southern emu-wren (*Stipiturus malachurus*). *Aust J Zool* 52: 583–604.
- Malan G (2004) The relative influence of prey abundance and co-breeders on the reproductive performance of polyandrous pale chanting-goshawks. *Ostrich* 75: 44–51.
- Manica LT, Marini MÂ (2012) Helpers at the nest of white-banded tanager Neothraupis fasciata benefit male breeders but do not increase reproductive success. J Ornithol 153: 149–159.
- Marzluff JM, Balda RP (1990) Pinyon jays: making the best of a bad situation by helping. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 197–237.
- Nakamura M (1998) Multiple mating and cooperative breeding in polygynandrous alpine accentors. I. Competition among females. *Anim Behav* 55: 259–275.
- Nias RC, Ford HA (1992) The influence of group-size and habitat on reproductive success in the superb fairy-wren *Malurus cyaneus*. *Emu* 92: 238–243.
- Noske RA (1991) A demographic comparison of cooperatively breeding and non-cooperative treecreepers (Climacteridae). *Emu* 91: 73–86.
- Noske RA (1998) Social organisation and nesting biology of the cooperatively-breeding varied sitella *Daphoenositta chrysoptera* in northeastern New South Wales. *Emu* 98: 85–96.

- O'brien TG (1997) Behavioural ecology of the North Sulawesi Tarictic hornbill *Penelopides exarhatus exarhatus* during the breeding season. *Ibis* 139: 97–101.
- Ocampo D, et al. (2012) Breeding biology of the red-bellied grackle (*Hypopyrrhus pyrohypogaster*): a cooperative breeder of the Colombian Andes. *Wilson J Ornithol* 124: 538–546.
- Patch-Highfill LD (2008) Estimating genetic diversity of Palila (*Loxioides bailleui*) and familial relationships of helper males. MSc thesis (Univ of Hawaii, Honolulu, HI).
- Pãdmaa T, Montgomerie R, Boag PT (1995) Mating system of the cooperatively breeding noisy miner *Manorina melanocephala*, as revealed by DNA profiling. *Behav Ecol Sociobiol* 37: 137–143.
- Rabenold PP, Rabenold KN, Piper WH, Haydock J, Zack SW (1990) Shared paternity revealed by genetic analysis in cooperatively breeding tropical wrens. *Nature* 348: 538–540.
- Radford JQ (2004) Breeding biology, adult survival and territoriality of the white-browed treecreeper (*Climacteris affinis*) in north-west Victoria, Australia. *Emu* 104: 305–316.
- Raitt RJ, Winterstein SR, Hardy JW (1984) Structure and dynamics of communal groups in the beechey jay. *Wilson Bull* 96: 206–227.
- Restrepo C, Mondrag ón ML (1998) Cooperative breeding in the frugivorous toucan barbet (*Semnornis ramphastinus*). *Auk* 115: 4–15.
- Reyer HU (1990) Pied kingfishers: ecological causes and reproductive consequences of cooperative breeding. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 529–557.
- Richardson DS, Burke T, Komdeur J (2002) Direct benefits and the evolution of female-biased cooperative breeding in Seychelles warblers. *Evolution* 56: 2313–2321.
- Ridgely RS, Robbins MB (1988) *Pyrrhura orcesi*, a new parakeet from southwestern Ecuador, with systematic notes on the *P. melanura* complex. *Wilson Bull* 100: 173–182.
- Ridley AR, Raihani NJ (2008) Task partitioning increases reproductive output in a cooperative bird. *Behav Ecol* 19: 1136–1142.
- Riehl C, Jara L (2009) Natural history and reproductive biology of the communally breeding greater ani (*Crotophaga major*) at Gat ún Lake, Panama. *Wilson J Ornithol* 121: 679–687.
- Rowley I (1978) Communal activities among white-winged choughs *Corcorax melanorhamphus*. *Ibis* 120: 178–97.
- Rowley I, Russell E (1995) The breeding biology of the white-winged fairy-wren *Malurus leucopterus leuconotus* in a western Australian coastal heathland. *Emu* 95: 175–184.
- Rowley I (1965) The life history of the superb blue wren, *Malurus cyaneus*. *Emu* 64: 251–297.
- Rubenstein DR (2007) Female extrapair mate choice in a cooperative breeder: trading

- sex for help and increasing offspring heterozygosity. *Proc R Soc B* 274: 1895–1903.
- Russell EM, Brown RJ, Brown MN (2004) Life history of the white-breasted robin, *Eopsaltria georgiana* (Petroicidae) in south-western Australia. *Aust J Zool* 52: 111–145.
- Russell E, Rowley I (2000) Demography and social organisation of the red-winged fairy-wren, *Malurus elegans*. *Aust J Zool* 48: 161–200.
- Russell E, Rowley I (1988) Helper contributions to reproductive success in the splendid fairy-wren (*Malurus splendens*). *Behav Ecol Sociobiol* 22: 131–140.
- Sankamethawee W, Pierce AJ, Hardesty BD, Gale GA (2011) Seasonal variability in survivorship of a cooperatively breeding tropical passerine. *Ecol Res* 26: 429–436.
- Saracura V, Macedo RH, Blomqvist D (2008) Genetic parentage and variable social structure in breeding Southern lapwings. *Condor* 110: 554–558.
- Seddon N, Tobias JA, Butchart SHM (2003) Group living, breeding behaviour and territoriality in the subdesert mesite. *Ibis* 145: 277–294.
- Shaw P, Owoyesigire N, Ngabirano S, Ebbutt D (2015) Life history traits associated with low annual fecundity in a central African Parid: the stripe-breasted tit *Parus fasciiventer*. *J Orithol* 156: 209–221.
- Sherley GH (1990) Cooperative breeding in riflemen (*Acanthisitta chloris*): benefits to parents, offspring and helpers. *Behaviour* 112: 1–22.
- Sherman PT (1995) Social organisation of cooperatively polyandrous white-winged trumpeters (*Psophia leucoptera*). *Auk* 112: 296–309.
- Sloane SA (1996) Incidence and origins of supernumeraries at bushtit (*Psaltriparus minimus*) nests. *Auk* 113: 757–770.
- Sridhar S, Karanth KP (1993) Helpers in cooperatively breeding small green bee-eaters (*Merops orientalis*). *Curr Sci* 65: 489–490.
- Strahl SD (1988) The social organization and behaviour of the hoatzin *Opisthocomus hoaxin* in central Venezuela. *Ibis* 130: 483–502.
- Sydeman WJ, Guntert M, Balda RP (1988) Annual reproductive yield in the cooperative pygmy nuthatch (*Sitta pygmaea*). *Auk* 105: 70–77.
- Tella JL (1993) Polyandrous trios in a population of Egyptian vultures (*Neophron percnopterus*). *J Raptor Res* 27: 119–120.
- Temple HJ, Hoffman JI, Amos W (2009) Group structure, mating system and extra-group paternity in the cooperatively breeding white-breasted thrasher *Ramphocinclus brachyurus*. *Ibis* 151: 99–112.
- Theuerkauf J, Rouys S, Mériot JM, Gula R, Kuehn R (2009) Cooperative breeding, mate guarding, and nest sharing in two parrot species of New Caledonia. *J Ornithol* 150: 791–797.
- Tingay RE, Watson RT (2002) Subordinate males sire offspring in madagascar fish-eagle (*Haliaeetus voc1feroes*) polyandrous breeding groups. *J Raptor Res* 36: 280–286.
- Uejima AMK, Boesing AL, Anjos LD (2012) Breeding and foraging variation of the plush-crested jay (*Cyanocorax chrysops*) in the Brazilian Atlantic forest. *Wilson*

- J Ornithol 124: 7-95.
- Valencia J, Cruz CDL, Gonz aez B (2003) Flexible helping behaviour in the azure-winged magpie. *Ethology* 109: 545–558.
- Varian-Ramos CW, Karubian J, Talbott V, Tapia I, Webster MS (2010) Offspring sex ratios reflect lack of repayment by auxiliary males in a cooperatively breeding passerine. *Behav Ecol Sociobiol* 64: 967–977.
- Wang SF (2011) Studies on the breeding ecology and social behavior of the masked laughingthrush (*Garrulax perspicillatus*). BA thesis (Sichuan Univ, Chengdu, China).
- Wilkinson R (1982) Social organization and communal breeding in the chestnut-bellied starling (*Spreo pulcher*). *Anim Behav* 30: 1118–1128.
- Williams DA, Rabenold KN (2005) Male-biased dispersal, female philopatry, and routes to fitness in a social corvid. *J Anim Ecol* 74: 150–159.
- Winterbottom M, Burke T, Birkhead TR (2001) The phalloid organ, orgasm and sperm competition in a polygynandrous bird: the red-billed buffalo weaver (*Bubalornis niger*). *Behav Ecol Sociobiol* 50: 474–482.
- Woolfenden GE, Fitzpatrick JW (1990) Florida scrub jays: a synopsis after 18 years of study. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 239–266.
- Woxvold IA, Mulder RA, Magrath MJL (2006) Contributions to care vary with age, sex, breeding status and group size in the cooperatively breeding apostlebird. *Anim Behav* 72: 63–73.
- Xu Y, Yang N, Zhang K, Yue B, Ran J (2011) Cooperative breeding by buff-throated partridge Tetraophasis szechenyii: a case in the Galliformes. *J Ornithol* 152: 695–700.
- Yuan HW, Liu M, Shen SF (2004) Joint nesting in Taiwan yuhinas: a rare passerine case. *Condor* 106: 862–872.
- Zack S (1986) Behaviour and breeding biology of the cooperatively breeding Grey-backed Fiscal Shrike *Lanius excubitorius* in Kenya. *Ibis* 128: 214–233.
- Zahavi A (1990) Arabian babblers: the quest for social status in a cooperative breeder. *Cooperative Breeding in Birds: Long-term Studies of Ecology and Behavior*, eds Stacey PB, Koenig WD (Cambridge Univ Press, Cambridge), pp 103–130.

Mammals

- Ågren G, Zhou Q, Zhong W (1989) Ecology and social behaviour of Mongolian gerbils, *Meriones unguiculatus*, at Xilinhot, Inner Mongolia, China. *Anim Behav* 37: 11–27.
- Bekoff M, Wells MC (1986) Social ecology and behavior of coyotes *Canus latrans*. *Advan Study Behav* 16: 251–338.
- Campbell RD, Rosell F, Nolet BA, Dijkstra VA (2005) Territory and group sizes in Eurasian beavers (*Castor fiber*): echoes of settlement and reproduction? *Behav Ecol Sociobiol* 58: 597–607.
- Cant MA (2000) Social control of reproduction in banded mongooses. Anim Behav 59:

- 147-158.
- Decanini DP, Macedo RH (2008) Sociality in *Callithrix penicillata*: II. Individual strategies during intergroup encounters. *Int J Primatol* 29: 627.
- Dietz JM, Baker AJ (1993) Polygyny and female reproductive success in golden lion tamarins, *Leontopithecus rosalia*. *Anim Behav* 46: 1067–1078.
- Doolan SP, Macdonald DW (1997) Band structure and failures of reproductive suppression in a cooperatively breeding carnivore, the slender-tailed meerkat (*Suricata suricatta*). *Behaviour* 134: 827–48.
- FitzGerald RW, Madison DM (1983) Social organization of a free-ranging population of pine voles, *Microtus pinetorum*. *Behav Ecol Sociobiol* 13: 183–187.
- Fuller TK, et al. (1992) Population dynamics of African wild dogs. *Wildlife 2001:* populations, eds McCullough DR, Barrett RH (Springer, Netherlands), pp 1125–1139.
- Garber PA, Moya L, Malaga C (1984) A preliminary field study of the moustached tamarin monkey (*Saguinus mystax*) in northeastern Peru: questions concerned with the evolution of a communal breeding system. *Folia Primatol* 42: 17–32.
- Getz LL, McGuire B, Pizzuto T, Hoffman J, Frase B (1993) Social organization of the prairie vole (*Microtus ochrogaster*). *J Mammal* 74: 44–58.
- Harrington FH, Mech DL, Fritts SH (1983) Pack size and wolf pup survival: their relationship under varying ecological conditions. *Behav Ecol Sociobiol* 13: 19–26.
- Jarvis JU (1981) Eusociality in a mammal: cooperative breeding in naked mole-rat colonies. *Science* 212: 571–573.
- Moehlman PD (1987) Social Organization in Jackals: The complex social system of jackals allows the successful rearing of very dependent young. *Am Sci* 75: 366–375.
- Moehlman PD (1979) Jackal helpers and pup survival. Nature 277: 382–383.
- Price EC (1992) The benefits of helpers: effects of group and litter size on infant care in tamarins (*Saguinus oedipus*). *Am J Primatol* 26: 179–190.
- Rood J (1990) Group size survival reproduction and routes to breeding in dwarf mongooses. *Anim Behav* 39: 566–572.
- Scharff A, Locker-Grütjen O, Kawalika M, Burda H (2001) Natural history of the giant mole-rat, *Cryptomys mechowi* (Rodentia: Bathyergidae), from Zambia. *J Mammal* 82: 1003–1015.
- Schradin C, Pillay N (2004) The striped mouse (*Rhabdomys pumilio*) from the succulent karoo of South Africa: a territorial group living forager with communal breeding and helpers at the nest. *J Comp Psychol* 118: 37–47.
- Smith MH (1966) The evolutionary significance of certain behavioral, physiological, and morphological adaptations of the old-field mouse, *Peromyscus polionotus*. PhD thesis (Univ of Florida, Gainesville, FL).
- Spinks AC, Bennett NC, Jarvis JUM (2000) A comparison of ecology of two populations of common mole rat, *Cryptomys hottentotus* the effect of aridity on food, foraging and body mass. *Oecologia* 125: 341–349.
- Svendsen GE (1980) Population parameters and colony composition of beaver

- (Castor canadensis) in Southeast Ohio. Am Midl Nat 104: 47–56.
- Thorington RW (1968) Observations on the tamarin *Saguinus midas*. *Folia Primatol* 9: 95–98.
- Wolff JO (1994) Reproductive success of solitarily and communally nesting white-footed mice and deer mice. *Behav Ecol* 5: 206–209.
- Ximenes MFFM, Sousa MBC (1996) Family composition and the characteristics of parental care during the nursing phase of captive common marmosets (*Callithrix jacchus*). *Primates* 37: 175–183.
- Zubiri CS, Gottelli D (1995) Spatial organization in the Ethiopian wolf *Canis simensis*: large packs and small stable home ranges. *J Zool* 237: 65–81.

Table S3. A preliminary summary of altruistic behaviors in the context of non-alloparenting in natural populations

| Type | Description of altruists' behavior | Taxon and sources |
|-------------------|--|---|
| Cooperative | Aggregate in lek to facilitate mating | Prairie mole cricket Gryllotalpa major ¹ , |
| courtship | success of related males at the expense of | Black grouse <i>Tetrao tetrix</i> ² , Wild turkey |
| | sacrificing their own mating opportunities | Meleagris galloparvo ³ , Peafowl Pavo |
| | | cristatus ⁴ , some manakins ^{5, 6, 7, 8, 9} |
| Egg dumping | Undertake the additional costs of parental | Lace bug Gargaphia solani ¹⁰ , some |
| | care to increase reproductive success of kin | waterfowl species11 |
| | parasites | |
| Anti-predatory or | Release costly chemical secretion to alert | Social aphids ¹² |
| parasite alarm | related individuals | |
| pheromone | | |
| Anti-predatory | Usually act as sentinels to protect kin | Some birds and mammals ¹³ |
| guard | against predation | |
| Killing parasite | Kill oppressors' offspring to reduce raiding | Social ants Temnothorax spp. 14, 15 |
| offspring | pressure on related host colonies nearby | |
| Adaptive suicide | Increase the probability of dying by | Pea aphid Acyrthosiphon pisum ¹⁶ |
| | dropping off plants to reduce the chance of | |
| | kin to be parasitized | |
| Larva cannibalism | Abandon to eat related larva but eat those | Tiger salamander Ambystoma tigrinum ¹⁷ , |
| | least related | Spadefoot toad Spea spp. 18 |

1. Keane *et al.* 2016; 2. Lebigre *et al.* 2014; 3. Krakauer 2005; 4. Petrie *et al.* 1999; 5. Shorey *et al.* 2000; 6. Loiselle *et al.* 2006; 7. McDonald & Potts 1994; 8. DuVal 2007; 9. Concannon *et al.* 2012; 10. Tallamy 2005; 11. Eadie & Lyon 2011; 12. Mondor & Messing 2007; 13. Caro 2005; 14. Pamminger *et al.* 2014; 15. Czechowski & Godzinska 2015; 16. McAllister & Roitberg 1987; 17. Pfennig & Collins 1993; 18. Pfennig & Frankino 1997.

Caro TM (2005) Antipredator Defenses in Birds and Mammals. (Univ of Chicago Press, Chicago).

Concannon MR, Stein AC, Uy JAC (2012) Kin selection may contribute to lek evolution and trait introgression across an avian hybrid zone. *Mol Ecol* 21: 1477-1486.

Czechowski W, Godzinska EJ (2015) Enslaved ants: not as helpless as they were thought to be. *Insectes Soc* 62: 9–22.

DuVal EH (2007) Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. *Am Nat* 169:423–432.

Eadie JM, Lyon BE (2011) The relative role of relatives in conspecific brood parasitism. *Mol Ecol* 20: 5114–5118.

Keane KT, Hill PSM, Booth W (2016) The kin selection hypothesis in a lekking mole cricket: assessing nested patterns of relatedness. *Biol J Lin Soc* 118: 382–393.

Krakauer AH (2005) Kin selection and cooperative courtship in wild turkeys. Nature

- 434: 69–72.
- Lebigre C, Alatalo RV, Soulsbury CD, Höglund J, Siitari H (2014) Limited indirect fitness benefits of male group membership in a lekking species. *Mol Ecol* 23: 5356–5365.
- Loiselle BA, et al. (2006) Kin selection does not explain male aggregation at leks of 4 manakin species. *Behav Ecol* 18: 287-291.
- McAllister MK, Roitberg BD (1987) Adaptive suicidal behaviour in pea aphids. *Nature* 328: 797–799.
- McDonald DB, Potts WK (1994) Cooperative display and relatedness among males in a lek-mating bird. *Science* 266: 1030–1032.
- Mondor EB, Messing RH (2007) Direct vs. inclusive fitness in the evolution of aphid cornicle length. *J Evol Biol* 20: 807–812.
- Pamminger T, Foitzik S, Metzler D, Pennings PS (2014) Oh sister where art thou? Spatial population structure and the evolution of an altruistic defence trait. *J Evol Biol* 27: 2443–2456.
- Petrie M, Krupa A, Burke T (1999) Peacocks lek with relatives even in the absence of social and environmental cues. *Nature* 401: 155–157.
- Pfennig DW, Collins JP (1993) Kinship affects morphogenesis in cannibalistic salamanders. *Nature* 362: 836–838.
- Pfennig DW, Frankino WA (1997) Kin-mediated morphogenesis in facultatively cannibalistic tadpoles. *Evolution* 51: 1993–1999.
- Shorey L, Piertney S, Stone J, Hoglund J (2000) Fine-scale genetic structuring on *Manacus manacus* leks. *Nature* 408:352–353.
- Tallamy DW (2005) Egg dumping in insects. Annu Rev Entomol 50: 347–370.